

On Propagator of Notoph

M. Bakalarska, W. Tybor*

Department of Theoretical Physics

University of Łódź

ul. Pomorska 149/153, 90-236 Łódź, Poland

Abstract

It is shown that the notoph propagator in the noncovariant longitudinal gauge is equivalent to the covariant Feynmann - like propagator.

*Supported by Łódź University Grant No. 505/581

1. Let us consider the interaction of the notoph [1] (i.e. the scalar particle described by the antisymmetric tensor field $B^{\mu\nu} = -B^{\nu\mu}$) with the external current $j^{\mu\nu} = -j^{\nu\mu}$. The Lagrangian is

$$\mathcal{L} = -\frac{1}{2}G_\mu G^\mu + \frac{1}{2}j_{\mu\nu}B^{\mu\nu}, \quad (1)$$

where $G^\mu = \partial_\nu B^{\nu\mu}$. The theory is invariant under the gauge transformation

$$\delta B^{\mu\nu} = \varepsilon^{\mu\nu\alpha\beta} \partial_\alpha \lambda_\beta, \quad (2)$$

(where λ^β is an arbitrary 4 - vector function), if the current obeys the conservation law

$$\varepsilon^{\mu\nu\alpha\beta} \partial_\nu j_{\alpha\beta} = 0. \quad (3)$$

The vector G^μ is invariant under the transformation (2) and it is a strength vector in the notoph theory.

The free action can be rewritten in the form

$$\int d^4x \left(-\frac{1}{2}G_\mu G^\mu \right) = \frac{1}{4} \int d^4x B_{\mu\nu} L^{\mu\nu\alpha\beta} B_{\alpha\beta}, \quad (4)$$

where the Lagrangian operator $L^{\mu\nu\alpha\beta}$ is

$$L^{\mu\nu\alpha\beta} = \square \Pi^{\mu\nu\alpha\beta} \quad (5)$$

and

$$\Pi^{\mu\nu\alpha\beta} = \frac{1}{2\square} \left(g^{\mu\alpha} \partial^\nu \partial^\beta + g^{\nu\beta} \partial^\mu \partial^\alpha - g^{\mu\beta} \partial^\nu \partial^\alpha - g^{\nu\alpha} \partial^\mu \partial^\beta \right). \quad (6)$$

The operator $\Pi^{\mu\nu\alpha\beta}$ has the symmetry properties of a Riemann tensor.

There is no inverse operator to the Lagrangian one because $\Pi^{\mu\nu\alpha\beta}$ is a projector operator

$$\Pi^{\mu\nu\alpha\beta} \Pi_{\alpha\beta}^{\sigma\lambda} = \Pi^{\mu\nu\sigma\lambda}.$$

The origin of the Lagrangian singularity is the gauge freedom of the theory. To fix the gauge we impose the gauge condition

$$\varepsilon^{\mu\nu\alpha\beta} \partial_\nu B_{\alpha\beta} = 0. \quad (7)$$

We note that this condition does not contradict the field equation when we switch on the interaction with the external current.

Let us introduce the effective free action

$$\int d^4x \left[-\frac{1}{2} G_\mu G^\mu + \frac{1}{8\alpha} (\varepsilon^{\mu\nu\sigma\lambda} \partial_\nu B_{\sigma\lambda})^2 \right] = \frac{1}{4} \int d^4x B_{\mu\nu} L_{(\alpha)}^{\mu\nu\sigma\lambda} B_{\sigma\lambda}, \quad (8)$$

where the Lagrangian operator

$$L_{(\alpha)}^{\mu\nu\sigma\lambda} = (1 - \frac{1}{\alpha}) \square \Pi^{\mu\nu\sigma\lambda} + \frac{1}{\alpha} \square E^{\mu\nu\sigma\lambda} \quad (9)$$

can be inverted for any finite value of the parameter α . The tensor

$$E^{\mu\nu\alpha\beta} = \frac{1}{2} (g^{\mu\alpha} g^{\nu\beta} - g^{\mu\beta} g^{\nu\alpha}) \quad (10)$$

is the unit tensor in a space of antisymmetric tensors.

The notoph propagator $D^{\mu\nu\alpha\beta}$ in the momentum space is defined as the inverse matrix to the Lagrangian one

$$L^{\mu\nu}_{\alpha\beta}(k, \alpha) D^{\alpha\beta\sigma\lambda} = E^{\mu\nu\sigma\lambda}. \quad (11)$$

The general form of the propagator is

$$D^{\mu\nu\alpha\beta} = \frac{A}{k^2} E^{\mu\nu\alpha\beta} + \frac{B}{k^2} \Pi^{\mu\nu\alpha\beta}(k). \quad (12)$$

From Eq. (11) we obtain

$$A = -\alpha, \quad B = \alpha - 1.$$

So, the notoph propagator is

$$D^{\mu\nu\alpha\beta} = -\frac{\alpha}{k^2} E^{\mu\nu\alpha\beta} + \frac{\alpha - 1}{k^2} \Pi^{\mu\nu\alpha\beta}(k). \quad (13)$$

If $\alpha = 0$, we get the propagator in the gauge (7). If $\alpha = 1$, we get the Feynmann - like propagator

$$D^{\mu\nu\alpha\beta}(k) = -\frac{1}{k^2} E^{\mu\nu\alpha\beta}. \quad (14)$$

We note that due to the current conservation, the effective propagator is the Feynmann - like one. Indeed, Eq. (3) can be rewritten in the form

$$\partial_\nu j_{\alpha\beta} + \partial_\alpha j_{\beta\nu} + \partial_\beta j_{\nu\alpha} = 0$$

and we get

$$\square j^{\alpha\beta} = \partial^\alpha j^\beta - \partial^\beta j^\alpha \quad (15)$$

where $j^\beta \equiv \partial_\nu j^{\nu\beta}$. With the help of Eqs (13) and (15) we obtain

$$D^{\mu\nu\alpha\beta} j_{\alpha\beta} = -\frac{\alpha}{k^2} j^{\mu\nu} + \frac{\alpha-1}{k^2} j^{\mu\nu} = -\frac{1}{k^2} j^{\mu\nu}.$$

The result is identical if the propagator (14) is used from the very beginning.

2. In Ref. [2] we have investigated the notoph theory in the (noncovariant) longitudinal gauge. We recall the main result of this paper.

In some fixed Lorentz frame we introduce the 3 - dimension notation. The decomposition of the notoph field is $B^{\mu\nu} = (\vec{E}, \vec{H})$, where

$$B^{k0} \rightarrow E_k, \quad B^{kn} \rightarrow \varepsilon_{knj} H_j, \quad k, n, j = 1, 2, 3.$$

The gauge freedom is removed completely by imposing the longitudinal gauge condition

$$rot \vec{E} = 0.$$

In this gauge the propagators of the notoph are

$$\begin{aligned} D_{ij}^{(E)}(k) &= \frac{k_i k_j}{|\vec{k}|^2} \frac{1}{k^2}; \\ D_{ij}^{(H)}(k) &= \left(\delta_{ij} - \frac{k_i k_j}{|\vec{k}|^2} \right) \frac{1}{|\vec{k}|^2}. \end{aligned} \quad (16)$$

In the Lorentz frame in that the gauge condition is imposed, we introduce the 4 - vector $\eta^\mu = (1, 0, 0, 0)$ [3]. Using this vector we can rewrite the propagators (16) in the 4 - dimension notation

$$\begin{aligned} D_{\mu\nu\alpha\beta}^{(E)}(k) &= \frac{1}{k^2} [(\eta k)^2 - k^2]^{-1} (\eta_\mu k_\nu - \eta_\nu k_\mu) (\eta_\alpha k_\beta - \eta_\beta k_\alpha); \\ D_{\mu\nu\alpha\beta}^{(H)}(k) &= -[(\eta k)^2 - k^2]^{-2} \{ g_{\mu\alpha} k_\nu k_\beta + g_{\nu\beta} k_\mu k_\alpha - g_{\mu\beta} k_\nu k_\alpha - g_{\nu\alpha} k_\mu k_\beta + \\ &\quad - (\eta_\mu \eta_\alpha k_\nu k_\beta + \eta_\nu \eta_\beta k_\mu k_\alpha - \eta_\mu \eta_\beta k_\nu k_\alpha - \eta_\nu \eta_\alpha k_\mu k_\beta) + \\ &\quad + (\eta k)^2 (g_{\mu\alpha} \eta_\nu \eta_\beta + g_{\nu\beta} \eta_\mu \eta_\alpha - g_{\mu\beta} \eta_\nu \eta_\alpha - g_{\nu\alpha} \eta_\mu \eta_\beta) + \\ &\quad - (\eta k) [g_{\mu\alpha} (k_\nu \eta_\beta + k_\beta \eta_\nu) + g_{\nu\beta} (k_\mu \eta_\alpha + k_\alpha \eta_\mu) + \\ &\quad - g_{\mu\beta} (k_\nu \eta_\alpha + k_\alpha \eta_\nu) - g_{\nu\alpha} (k_\mu \eta_\beta + k_\beta \eta_\mu)] \}. \end{aligned} \quad (17)$$

Let us consider the exchange of the notoph between two currents

$$A = \frac{1}{4} j^{\mu\nu}(-k) D_{\mu\nu\alpha\beta}(k, \eta) j^{\alpha\beta}(k) \quad (18)$$

where

$$D_{\mu\nu\alpha\beta}(k, \eta) = D_{\mu\nu\alpha\beta}^{(E)}(k) + D_{\mu\nu\alpha\beta}^{(H)}(k). \quad (19)$$

Using the current conservation law (3) we can rewrite the amplitude (18) with the help of the Feynmann - like propagator (14)

$$A = \frac{1}{2} j^{\mu\nu}(-k) D_{\mu\nu\alpha\beta}(k) j^{\alpha\beta}(k). \quad (20)$$

So, the notoph propagator in the longitudinal gauge is equivalent to the covariant Feynmann - like propagator.

3. Let us note that the covariant form (14) of the notoph propagator can be confirmed in another way. Performing the canonical analysis of the theory described by the Lagrangian (1), we get the physical Hamiltonian of the interaction

$$\mathcal{H}_{int} = -j_{2Lk} B_L^{0k} + \frac{1}{2} j_{1T}^i \frac{1}{\Delta} j_{1Ti},$$

or, in the momentum space

$$\mathcal{H}_{int} = -j_{2Lk}(-k) B_L^{0k}(k) - \frac{1}{2} |\vec{k}|^{-2} j_{1T}^i(-k) j_{1Ti}(k) \quad (21)$$

where

$$j_1^i \equiv j^{0i}, \quad j_2^i = -\frac{1}{2} \varepsilon^{imn} j_{mn},$$

and the mark $L(T)$ denotes the longitudinal (transversal) part of the 3 - vector. Using the standard methods of the S - matrix formalism [3], with the help of this Hamiltonian, we can calculate the amplitude of the exchange of the gauge field between two external currents. We get

$$A = -\frac{j_{2Li}(-k) j_{2L}^i(k)}{k^2} - \frac{j_{1Ti}(-k) j_{1T}^i(k)}{|\vec{k}|^2}.$$

It is the amplitude (20) rewritten in the component form with the help of the current conservation law (3).

4. Let us consider in es detail the case of the strong conserving current [4]

$$j^{\mu\nu} = \partial^\mu J^\nu - \partial^\nu J^\mu, \quad (22)$$

where J^μ is a 4 - vector (or a 4 - pseudovector). In this case the amplitude (20) is

$$A = J^\mu(-k) D_{\mu\nu}(k) J^\nu(k), \quad (23)$$

where (see Ref. [1])

$$D_{\mu\nu}(k) = g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}. \quad (24)$$

To understand this result we observe that from the field equation

$$\partial_\mu G_\nu - \partial_\nu G_\mu = -\partial_\mu J_\nu + \partial_\nu J_\mu$$

we get

$$G^\mu = -J^\mu - \partial^\mu \varphi,$$

where $\varphi(x)$ is a scalar field. The interaction term $\frac{1}{2} j^{\mu\nu} B_{\mu\nu}$ with the current (22) is equivalent to

$$-J_\mu G^\mu = J_\mu J^\mu + \partial_\mu \varphi J^\mu,$$

what is in agreement with Eqs (23) and (24). So, the notoph is propagated if the current J^μ is not conserved. The process of the notoph radiation (or absorption) is possible too, if the current does not conserve. Indeed, such processes are described by the vertex $J_\mu \partial^\mu \varphi$.

Interaction of the notoph with other fields (via the strong conserving current) have been discussed in Ref. [1]. To illustrate the main features of such interactions we consider, for example, the interaction with the Dirac field ψ

$$\mathcal{L} = -\frac{1}{2} G_\mu G^\mu - J_\mu G^\mu + \bar{\psi}(-i\gamma_\mu \partial^\mu + m)\psi$$

where $J^\mu = \bar{\psi}\gamma^\mu\psi$ or $J^\mu = \bar{\psi}\gamma^\mu\gamma_5\psi$. In the first case the notoph is not radiated and propagated: the local $J^\mu J_\mu$ interaction appears. In the second case the notoph can be radiated and propagated: the local term $J^\mu J_\mu$ appears too.

We are grateful to Prof. J. Rembieliński and Dr B. Broda for interesting discussions.

References

- [1] V. Ogievetsky, I. Polubarinov, *Yad. Fiz.* **4**, 216 (1966).
- [2] L. C. Papaloucas, J. Rembieliński, W. Tybor, *Acta Phys. Pol.* **B22**, 429 (1991).
- [3] J. D. Bjorken, S. D. Drell, *Relatywistyczna teoria kwantów, (Relativistic Quantum Mechanics, Relativistic Quantum Fields)*, PWN, Warszawa (1985), pp. 430 - 433.
- [4] J. N. Goldberg, *Phys. Rev.* **89**, 263 (1953).